

Durham Research Online

Deposited in DRO:

06 April 2016

Version of attached file:

Accepted Version

Peer-review status of attached file:

Peer-reviewed

Citation for published item:

Racionero-Gómez, B. and Sproson, A.D. and Selby, D. and Gröcke, D.R. and Redden, H. and Greenwell, H.C. (2016) 'Rhenium uptake and distribution in phaeophyceae macroalgae, *Fucus vesiculosus*.', *Royal Society open science.*, 3 (5). p. 160161.

Further information on publisher's website:

<http://dx.doi.org/10.1098/rsos.160161>

Publisher's copyright statement:

© 2016 The Authors. Published by the Royal Society under the terms of the Creative Commons Attribution License <http://creativecommons.org/licenses/by/4.0/>, which permits unrestricted use, provided the original author and source are credited.

Additional information:

Use policy

The full-text may be used and/or reproduced, and given to third parties in any format or medium, without prior permission or charge, for personal research or study, educational, or not-for-profit purposes provided that:

- a full bibliographic reference is made to the original source
- a [link](#) is made to the metadata record in DRO
- the full-text is not changed in any way

The full-text must not be sold in any format or medium without the formal permission of the copyright holders.

Please consult the [full DRO policy](#) for further details.

ROYAL SOCIETY
OPEN SCIENCE

**Rhenium uptake and distribution in Phaeophyceae
macroalgae, *Fucus vesiculosus***

Journal:	<i>Royal Society Open Science</i>
Manuscript ID	RSOS-160161.R1
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Racionero-Gómez, Blanca; Durham, Sproson, Adam; Durham University, Selby, Dave; University of Durham, Earth Sciences Grocke, Darren; Durham University Redden, Hilary; Durham University, Greenwell, Christopher; Durham University, Chemistry
Subject:	biochemistry < CROSS-DISCIPLINARY SCIENCES, biogeochemistry < CROSS-DISCIPLINARY SCIENCES, Chemical biology < CHEMISTRY
Keywords:	rhenium, macroalgae, uptake, distribution, phytomining, bioremediation
Subject Category:	Biology (whole organism)

SCHOLARONE™
Manuscripts

Rhenium uptake and distribution in Phaeophyceae macroalgae, *Fucus vesiculosus*
B. Racionero-Gómez,¹ A. D. Sproson¹, D. Selby¹, D. R. Gröcke¹, H. Redden^{1,2} & H. C. Greenwell^{1,2}

[1] {Department of Earth Sciences, Durham University, Durham, DH1 3LE, UK}
[2] {Department of Chemistry, Durham University, Durham, DH1 3LE, UK}
Correspondence to: B. Racionero Gómez (blancaraci@gmail.com telephone: +34646976656) and H. C. Greenwell (chris.greenwell@durham.ac.uk telephone: +441913342324 fax: +44 0191 3342301)

Abstract

Owing to Re having no known biological role, it is not fully understood how Re is concentrated in oil kerogens. A commonly held assumption is that Re is incorporated into decomposing biomass under reducing conditions. However, living macroalgae also concentrates Re to several orders of magnitude greater than that of seawater. This study utilizes *Fucus vesiculosus* to assess Re uptake and its subsequent localisation in the biomass. It is demonstrated that the Re abundance varies within the macroalgae and that Re is not located in one specific structure. In *F. vesiculosus* the uptake and tolerance of Re was evaluated via tip cultures grown in seawater of different Re(VII) compound concentrations (0 to 7450 ng/g). A positive correlation is shown between the concentration of Re doped seawater and the abundance of Re accumulated in the tips. However, significant differences between Re(VII) compounds are observed. Although the specific cell structures where the Re is localised is not known, our findings suggest that Re is not held within chloroplasts or cytoplasmic proteins. In addition, metabolically inactivated *F. vesiculosus* does not accumulate Re, which indicates that Re uptake is via syn-life bioadsorption/bioaccumulation and that macroalgae may provide a source for Re phytomining and/or bioremediation.

Keywords

Rhenium, macroalgae, uptake, distribution, phytomining, bioremediation

Acknowledgements

We wish to express special thanks to Dr Carl Patterson, Dr Deenah Osman and Prof. Nigel Robinson for their help with the protein column preparation, protein analysis and discussions, and to Dr Chris Ottley, Dr Joanna Hesselink and Dr Emily Unsworth for their help with the ICP-MS. We are very grateful to OEA Labs Ltd for funding the project. We thank the school of Biological and Biomedical Sciences of Durham University for the culture experiment facilities. Finally, we acknowledge the constructive remarks by Bernhard Peucker-Ehrenbrink and an anonymous reviewer, which have improved this manuscript.

Introduction

The behaviour of Rhenium (Re) in seawater is defined by the low reactivity of the perrhenate ion (ReO_4^- ; Re(VII)), which is the only significant Re species found in ocean waters [1]. The concentration of Re in the open ocean (0.009 – 0.0074 ng/g; [2,3]) is a factor of three higher than average river water (~0.005 pg/g; [4]) and much lower compared to terrestrial environments (continental crust values of 0.2 – 2 ng/g; organic-rich sedimentary rocks values 0.2 – 100 ng/g; [5] and references therein) and sulphide minerals (low ng/g to hundreds of mg/g; [6]).

Although the Re concentration in seawater is low in comparison to the terrestrial realm and despite there being no known biological use of Re, marine macroalgae (i.e. seaweed), especially brown macroalgae, are known to concentrate Re up to several hundreds of ng/g [7–9], in addition to many metal cations and oxoanions through forming a variety of metal complexes with, for example, alginate, proteins, polysaccharides of the cell wall, fucans, etc. [10]. To date, positively charged metals associated with macroalgae have been extensively

studied [11–14], however, relatively little is known about the mechanisms by which macroalgae uptake negatively charged metal oxoanions such as the perrhenate ion. Experiments have shown that Re is most likely stored within algal cells, rather than on the algal cell surface or within the intercellular matrix [9,15]. Specifically, it has been proposed that protonated amino groups could be involved, forming an ion pair with perrhenate [15,16]. Moreover, Kim *et al.* [17] showed that ReO_4^- interacted strongly with chitosan, a cationic polymer of glucosamine. Chitosan is only reported in nature in some fungi, crustacea and the termite queen's abdominal wall. However, Nishino *et al.*, [18] isolated and characterized a novel polysaccharide containing an appreciable amount of glucosamine in *F. vesiculosus*, which suggests a further route to possible Re uptake.

Assuming that Re is being stored inside the macroalgae cells, a mechanism for Re uptake into the cells should be identifiable. Macroalgae could inadvertently take up ReO_4^- (ionic radius of 2.60 Å) by confusing it for phosphate (PO_4^{3-}) (ionic radius of 2.38 Å). A similar mechanism is proposed for arsenate (AsO_4^{3-}) [20]. Sulphate (SO_4^{2-}), nitrate (NO_3^-) and Chloride (Cl^-) also have similar ionic radius to ReO_4^- (i.e. 2.58 Å, 1.96 Å and 1.81 Å, respectively). Thus these ions could be also competing with ReO_4^- . For instance [19] showed that there is a positive correlation between K^+ and technetium (Tc) accumulated in three plant species (*Cucumis sativus L.*, *Raphanus sativus L.* and *Brassica chinensis L.*) and explained this as a result of TcO_4^- being taken up by mistaken identity for Cl^- , as a counter ion for K^+ uptake. As Re is a Tc analogue [17, 9, 21], ReO_4^- might be taken up in a similar manner. In addition, competitive incorporation between ReO_4^- and NO_3^- in sodalites has also been found [22], however as sodalite is a mineral ReO_4^- incorporation cannot be compared with ReO_4^- concentration in biologically active organisms.

Importantly, understanding the uptake of Re will help to elucidate the uptake of Tc, which is produced in nuclear power stations. Moreover, a better knowledge on the uptake mechanism

could open the possibility to use macroalgae as bioconcentrators of Re and Tc, thus bioremediation of Tc contaminated waters and phytomining of Re could be achieved using *F. vesiculosus*, as well as potentially providing an alternative hypothesis for the high concentration of Re within oil forming kerogens.

This study uses a brown macroalgae (Phaeophyceae) to establish: (i) where Re is stored; (ii) the limit of Re uptake; and (iii) the uptake mechanism of Re (i.e. active concentration in which the transport requires energy to oppose the concentration gradient, or passive concentration, with transport requiring no energy and entirely correlated with the concentration). The Re abundance data for the different structures of *F. vesiculosus*: holdfast, stipe, fertile tips, non-fertile tips, vesicles and blades (Fig. 1), and isolated cytoplasmic proteins and chloroplasts is investigated. The uptake limit of Re in macroalgae is determined via cultures of *F. vesiculosus* under different ReO_4^- concentrations and using different ReO_4^- chemical compounds (i.e. HReO_4 (Re metal dissolved in HNO_3), KReO_4 , NaReO_4 and NH_4ReO_4). Cultured *versus* dead macroalgae were used to provide insight into the uptake mechanism of ReO_4^- by macroalgae.

Material and methods

- Macroalgae used in the study: *Fucus vesiculosus*

The available Re data for brown macroalgae (Phaeophyceae) indicate it has the highest Re accumulation of all macroalgae, with *Fucus vesiculosus* possessing the highest Re concentrations measured to date for a macroalgae [7]. *F. vesiculosus* is a common macroalgae found along sheltered shores of the North Sea, Baltic Sea, Atlantic Ocean and Pacific Ocean. *F. vesiculosus* is a tethered macroalgae with air bladders that are produced annually allowing the individual fronds to float. The growth rate ranges between 0.05 and 0.14 cm/day [23,24] and they have a life span in the order of 3 to 5 years [25]. The species is

1
2
3 98 annually episodic, gonochoristic and highly fecund (i.e. prolific) [25]. Gametes are released
4
5 99 into the seawater and the eggs are fertilized externally to form a zygote that starts to develop
6
7 100 as soon as it settles into a substrate [26]. The gametes are released from receptacles, which
8
9
10 101 are found in the fertile tips of the macroalgae. However, *F. vesiculosus* also has non-fertile
11
12 102 tips without these structures. Non-fertile tips are composed by a parenchymatous thallus (i.e.
13
14 103 tissue like structure) [25–27]. The structures of *F. vesiculosus* are shown in Fig. 1.
15
16

17 104 - **Macroalgae collection sites**
18
19

20 105 Five specimens of *F. vesiculosus* were collected from Staithes, North Yorkshire, UK
21
22 106 (54°33'N 00°47'W) in May, 2014. These samples were used to determine the Re abundance
23
24 107 of specific structures of the macroalgae. An additional six samples were collected each month
25
26 108 at Boulmer Beach, Northumberland, UK (55°25'N 1°34'W) in May, June, October and
27
28 109 November in 2014, and January to June in 2015, for fertile and non-fertile tip separation, all
29
30 110 the culture experiments, chloroplast isolation and protein purification.
31
32
33

34 111 - **Rhenium abundance and distribution in macroalgae structures**
35
36

37 112 Prior to analysis all specimens were kept individually in plastic sample bags for transport,
38
39 113 and stored in a freezer (-10 °C) for 48 h. Each specimen was washed and soaked in deionised
40
41 114 (Milli-Q™) water to remove any attached sediment and salt. To establish the abundance and
42
43 115 distribution of Re in the macroalgae the sample was divided into different structural
44
45 116 components; fertile tips, non-fertile tips, vesicles, stipe, holdfast, blades (Fig. 1). In addition,
46
47 117 all the algae components were mixed to assess an average Re abundance. Each structure was
48
49 118 dried in an oven at 60 °C for 12 h.
50
51
52

53
54 119 - **Rhenium uptake of macroalgae**
55
56
57
58
59
60

120 To investigate the uptake of Re by macroalgae, non-reproductive apical thallus tips of nine *F.*
121 *vesiculosus* specimens (length = > 1.5 cm; wet weight (WW) = 0.12–0.15 g), without visible
122 microalgae (i.e. epiphytes), from Boulmer beach were cultured in seawater (modified after
123 Gustow *et al.*; [28]) with a known concentration of Re. In brief, the culture experiments were
124 performed using a 250 mL glass jar containing two mesh shelves. Three tips were placed in
125 the bottom of the jar and three tips to each mesh, having in total nine tips, with each set of
126 tips taken from a different specimen (Fig. 2). All jars were filled with sterile filtered (0.7 µm)
127 seawater from Boulmer beach. A huge diversity of macroalgae grow naturally at Boulmer
128 beach, thus water obtained at Boulmer water is expected to be nutrient replete as it permits
129 the growth of a wide variety of species. Each set of three jar replicates were doped using a
130 known volume of ReO_4^- from different Re compounds: an already prepared solution of Re
131 metal with nitric acid (HReO_4) (i.e. 83787 Sigma Aldrich) or commercially obtained Re(VII)
132 salts (KReO_4 , NH_4ReO_4 and NaReO_4).

133 HNO_3 dissolves Re metal forming HReO_4 , [29]. For the cultures using HReO_4 , Boulmer
134 seawater ReO_4^- concentration was analysed. The Re abundance in the seawater was
135 determined by isotope dilution ICP-MS (details below). The seawater possess a Re
136 abundance of ~ 0.007 ng/g (6.95 ± 0.19 pg/g) coinciding with the concentrations reported by
137 Anbar *et al.* [2]. The seawater culture experiments were conducted in Re concentrations are
138 equal to that of seawater, and 10×, 50×, 100×, 500×, 1000×, 2667×, 10000×, 133333× and
139 266667× that of the concentration of seawater (i.e. 0.007 ng/g, 0.075 ng/g, 0.373 ng/g, 0.745
140 ng/g, 3.725 ng/g, 7.450 ng/g, 20 ng/g, 75 ng/g, 1000 ng/g and 2000 ng/g, respectively). In
141 addition, three jars were filled with artificial seawater that was not doped with Re, and one jar
142 was doped with a concentration a million times that of the Re seawater concentration in order
143 to reach an extreme concentration of 7450 ng/g.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

144 For the cultures using Re(VII) (perrhenate) salts, the same approach was used, where the
145 doped Re concentrations of seawater in the cultures were 10×, 50×, 100× and 1000× that of
146 seawater (i.e. 0.075 ng/g, 0.373 ng/g, 0.745 ng/g and 7.45 ng/g, respectively).

147 To reduce evaporation, while allowing gaseous exchange with the atmosphere, all the jars
148 were loosely covered with lids. No additional nutrients were added into the seawater or
149 artificial seawater. The algae tips inside the bottles were transferred into an incubator with a
150 set light/dark rhythm of 16:8, light intensity of 125 $\mu\text{mol photons/m}^2\cdot\text{s}^2$ and a temperature of
151 11°C. The wet weight (WW) of the algal tips, per jar, was measured every 2–3 days during 25
152 days of the culturing period for all cultures except the cultures of June 2015, which only
153 lasted 15 days. At the same time, the media was changed (between 4 and 7 times for all
154 cultures) to avoid accumulation of metabolites and replenish nutrients. The salinity (~35 ppt)
155 of the Re doped seawater did not appreciably change from that of natural seawater collected
156 from Boulmer and remained constant throughout the culture experiments. The pH (~9.0),
157 however, changed from that of the natural seawater collected from Boulmer (~8.2) due to the
158 metabolic activity of the macroalgae (photosynthesis) and remained constant throughout the
159 culture experiments.

160 Two additional sets of culture experiments were conducted to establish if ReO_4^- is taken up
161 by syn-life bioabsorption/bioaccumulation or passive processes. Understanding syn-life
162 bioaccumulation and bioabsorption as the biological sequestration of substances or chemicals
163 through any route at a higher concentration than that at which it occurs in the surrounding
164 environment/medium when macroalgae is metabolically active (i.e. alive) [30]. Therefore, in
165 order to assess bioaccumulation, non-reproductive thallus tips were killed through either
166 boiling, drying or freezing. Specifically, non-reproductive thallus tips ($n = 81$) from Boulmer
167 beach were heated for 2 h at 100 °C, and a further 21 tips were heated at 100 °C for only 5

168 min. Additionally, 21 non-reproductive thallus tips were air dried for 72 h and another 21 tips
169 were frozen with liquid nitrogen. In total, 18 jars were filled with sterile (i.e. autoclaved at
170 121°C for 30 min) and filtered (0.7 µm) seawater from Boulmer beach. The jars containing
171 boiled tips were divided into three subgroups composed of three replicates of each with the
172 following treatments: seawater and seawater doped with 7.45 ng/g of HReO₄. The other set of
173 three replicates containing dried, boiled (5 min) or frozen non-reproductive thallus tips,
174 respectively, were only treated with seawater spiked with 7.45 ng/g HReO₄.

175 In order to re-confirm the uptake mechanism, four tips were placed in the bottom of the jar
176 and four tips to each mesh, having in total 12 tips of different specimens in each jar. All jars
177 were filled with sterile filtered (0.7 µm) seawater from Boulmer beach and doped with 7.45
178 ng/g NaReO₄. After 3 days the media solution was changed and set to 0.075 ng/g of NaReO₄
179 and, finally, after another 3 days the media solution was changed and not doped. Prior to each
180 change of the media four sample tips were taken for Re analysis.

181 - **Chloroplast isolation**

182 A procedure modified from Popovic *et al.* [31] was used for the isolation of chloroplasts.
183 Approximately 10 g of non-reproductive thallus tips were cut into 2 mm² pieces using
184 scissors. These were washed by stirring with 2 L of filtered seawater with 75 mL of grinding
185 medium added. The grinding medium consisted of 1 M sorbitol, 2 mM MnCl₂, 1 mM MgCl₂,
186 0.5 mM K₂HPO₄, 5 mM EDTA, 2 mM NaNO₃, 2 mM ascorbate, 2 mM cysteine, 0.2 % (w/v)
187 BSA and 50 mM of MES Buffer (pH 6.1). All the subsequent steps were undertaken in ice
188 water. The washed tissue was divided into two portions, each ground with a mortar and
189 pestle, increasing gradually the volume to 50 mL. Then, each portion was diluted into 100
190 mL of medium and passed through a stainless steel strainer and four layers of cheese cloth.
191 Chloroplasts were isolated by centrifugation for 7 min at 5500 G. The pellet was re-

1
2
3 192 suspended with 10 mL of a reaction medium containing 1 M sorbitol, 1 mM MnCl₂, 1 mM
4
5 193 MgCl₂, 2 mM EDTA, 0.5 mM K₂HPO₄ and 50 mM HEPES (pH 8.1). Another centrifugation
6
7 194 at 5500 G for 7 min was performed and chloroplasts were re-suspended with 2 mL of HEPES
8
9 195 buffer. To test the isolation, the absorbance spectrum of the last solution obtained was
10
11 196 observed under a light microscope. The extracted chloroplasts were preserved using HEPES
12
13 197 (as it does not contain Re) and stored in a fridge for 3 days. In order to remove HEPES from
14
15 198 the chloroplasts the HEPES-chloroplast mixture was centrifuged. The chloroplast pellet was
16
17 199 white-brown and the HEPES solution was green-brown. The observation showed that the
18
19 200 pigments had released and were free in the solution.
20
21
22

23
24 201 - **Cytoplasmic proteins isolation**
25

26
27 202 A procedure modified from Boer *et al.* [32] was employed for the isolation of cytoplasmic
28
29 203 proteins. Approximately 2 g of freshly ground non-reproductive thallus tips were used for
30
31 204 protein extraction. The tips were mixed with 9 mL of 10 mM HEPES (pH 7.8) buffer,
32
33 205 vortexed and centrifuged twice at 1000 G for 1 min. The homogenate was sonicated for 1
34
35 206 min, 10 times and centrifuged at 4500 G for 5 min. The supernatant was centrifuged at 14000
36
37 207 G for 10 min. A 60 mM saturated CaCl₂ solution was used to re-suspend the pellet, which
38
39 208 was agitated and then centrifuged at 14000 G for 5 min. The supernatant was then separated
40
41 209 via gel filtration (i.e. size exclusion column chromatography). A PD-10 Desalting Column
42
43 210 containing *Sephadex G-25 Medium* as matrix was used to separate molecules from the
44
45 211 supernatant by their molecular size. Larger molecules than the *Sephadex* matrix pores are
46
47 212 eluted first and smaller molecules than the matrix pores are eluted later, depending on the
48
49 213 molecular size, the molecules will penetrate the matrix pores to varying extent. The
50
51 214 separation was carried out following the gravity protocol detailed in PD-10 Desalting Columns
52
53 215 Instructions [33] using the same buffer described above. 1 mL elution fractions obtained were
54
55
56
57
58
59
60

1
2
3 216 analysed by ICP-MS after being diluted 10 times with 0.8 N HNO₃. Protein content of the
4
5 217 fractions was analysed based on the absorbance shift of the dye Coomassie Brilliant Blue G-
6
7 218 250.
8
9

10
11 219 - **Re abundance determinations and data treatment**
12

13 220 Rhenium abundance determinations for all samples were obtained at the Durham
14
15 221 Geochemistry Centre in the Laboratory for Sulphide and Source Rock Geochronology and
16
17 222 Geochemistry. Each sample of *F. vesiculosus* was oven-dried at 60 °C for 24 h and ground
18
19 223 into a powder with an agate mortar and pestle. Approximately 100 mg of the sample powder
20
21 224 was spiked. Abundances were obtained by both direct calibration and isotope-dilution
22
23 225 methodologies (Table 1, 2, 3, 4 and 5). For the latter samples were doped with a known
24
25 226 amount of ¹⁸⁵Re tracer solution (isotope dilution methodology). The sample and if used, the
26
27 227 tracer solution, were digested in a mix of 3 ml of 12 N HCl and 6 ml of 16 N HNO₃ at 120 °C
28
29 228 overnight in a PFA Savillex 22 mL vial. The dissolved sample solution was evaporated to
30
31 229 dryness at 80 °C. The rhenium abundance of seawater from Boulmer beach was determined
32
33 230 by isotope dilution-ICP-MS. Approximately 30 mL of seawater was doped with a known
34
35 231 amount of ¹⁸⁵Re tracer solution and evaporated. The rhenium fraction was further purified
36
37 232 using standard anion chromatography methodology. Rhenium for all macroalage samples was
38
39 233 isolated from the dried sample using 5 mL 5 N NaOH 5 mL acetone solvent extraction
40
41 234 procedure [8,34]. The Re-bearing acetone was evaporated to dryness at 60 °C. For ICP-MS
42
43 235 the dried Re fraction was dissolved in 1.2 mL of 0.8 N HNO₃. For thermal ionization mass
44
45 236 spectrometry in negative ion mode (N-TIMS) analysis the purified Re fraction was loaded
46
47 237 onto a Ni wire filament, with the Re isotope compositions determined using Faraday cup
48
49 238 measurements on a Thermo Scientific TRITON mass spectrometer. Total procedural blanks
50
51 239 are 1 ± 0.1 pg (n = 6). For samples analysed by isotope dilution to determine absolute Re
52
53
54
55
56
57
58
59
60

1
2
3 240 abundance, all sources of uncertainty (e.g., standard measurement, isotope measurement,
4
5 241 calibration of the tracer solution, fractionation correction and blank values) are propagated to
6
7 242 yield a final uncertainty. For direct calibration, prior to each analysis, an instrument
8
9 243 performance check confirm satisfactory performance of the ICP-MS. Five freshly prepared
10
11 244 standards were made each time and formed calibration lines with an R value > 0.999 and <
12
13 245 2% RSD uncertainty. Moreover, all the samples had a reproducibility of < 5% RSD.
14
15
16
17 246 Statistical analysis, t-test and Tukey's HSD tests, using a significance level of 0.05, were
18
19 247 performed using R Studio software. For testing the statistical hypothesis, *p*-values are used.
20
21 248 The *p*-value is defined as the probability of obtaining a result more extreme or equal to what
22
23 249 was actually observed, thus, if *p*-value is smaller or equal to the significance level, it suggests
24
25 250 that the observed data are consistent with the hypotheses.

26
27
28
29 251 **Results**

30
31
32 252 - **Location of Re within *F. vesiculosus* structures**

33
34
35 253 All analyzed structures of *F. vesiculosus* are naturally enriched in Re by approximately one
36
37 254 thousand times that found in seawater (Fig. 1). The contents of Re range from 23 to 313 ng/g
38
39 255 (Fig. 1). Significant differences were observed (*p*-value: 0.02) between the five samples of
40
41 256 macroalgae tips (~126 ng/g) and the sample representing a mix of the plant components (~74
42
43 257 ng/g). Further, significant differences were also observed (*p*-value: 0.003) between fertile
44
45 258 (~123 ng/g) and non-fertile tips (~313 ng/g) (Fig. 1).

46
47
48
49 259 - **Uptake of Re by *F. vesiculosus* culture tips**

50
51
52 260 The natural Re abundance of the seawater collected from Boulmer beach and utilised for the
53
54 261 culture experiments is 6.95 ± 0.19 pg/g (~0.007 ng/g), which is in agreement with previous
55
56 262 studies of coastal waters [2]. The results shown in Figures 3, 4 and 5 indicate that in 25 days
57
58
59
60

the Re content of the macroalgae increased proportionally to the amount of Re species doped in the seawater. However, variation in the uptake capacity by *F. vesiculosus* of the different ReO_4^- compounds doped in seawater is observed. Moreover, a significant variation (p-value < 0.05) in uptake capacity between months of collection (i.e. February, March, May and June cultures with Re(VII) salts) was observed only after 0.37 ng/g of doped Re(VII) in the media. March cultures accumulated ~7000 ng/g more Re than February, May and June culture tips (Table 6). Moreover, cultures doped with HReO_4 and Re(VII) salts also show different amounts of accumulation. The accumulation of Re in *F. vesiculosus* grown with all Re(VII) salts is significantly lower (p-value < 0.05) than the accumulation obtained with cultures made with HReO_4 , also only after 0.37 ng/g of doped Re to the media (Fig. 3). It is observed that cultures in Re doped solution made from HReO_4 take up 50% of the amount of Re in seawater, in contrast to only 0.03–15% for solution doped with Re from Re(VII) salts (Table 6). Because of this, cultures with high concentrations of ReO_4 in the media were made only with HReO_4 . A linear correlation is observed between the amount of Re doped in the cultures and the accumulation of Re in the alive cultured macroalgae until an accumulation of 63284 ng/g of Re was reached, after which Re uptake ceased as the macroalgae died (Fig. 4). We also observed there is a limit on the uptake of Re in the cultured macroalage between 75 and 1000 ng/g of HReO_4 in the seawater media. Furthermore, visually the macroalgae tips grown in high concentrations (2000 and 7450 ng/g) did not seem as metabolically active as those in lower concentrations. In total, macroalgae tips extracted up to ~60000 ng/g of Re in 25 days (see Fig. 4 and 5).

F. vesiculosus non-fertile tips under 7.45 ng/g of NaReO_4 in the media, after 3 days were capable of accumulating ~150 ng/g more than the background Re concentration in them (Fig. 6). These tips were then transferred to subsequent lower concentrations of NaReO_4 (0.075 and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

0.007 ng/g) and exhibited accumulations of ~100 ng/g more than the background concentration of Re. Therefore a release of 50 ng/g was found after transference (Fig. 6).

In comparison to living organism samples, *F. vesiculosus* non-fertile thallus tips metabolically deactivated by boiling, freezing with liquid nitrogen or drying showed appreciably little to no accumulation of Re (between 36 and 19 ng/g) compared to the concentration reached in fresh tips (i.e. alive) (~16000 ng/g) with same HReO₄ concentrations in the media of 7.45 ng/g (see Fig. 7). Also, the majority of the Re content in the macroalgae was released in the media within the first 2–3 days of the experiment and the media turned brown.

- **Chloroplast isolation**

Chloroplasts were isolated from *F. vesiculosus* non-fertile tips. The non-fertile tips as a whole contain between 100 and 200 ng/g of Re. Chloroplasts are found throughout the whole macroalgae organism, although exist in greater abundance in the non-fertile tips. Both the HEPES solution and the chloroplast pellet were analysed. 1 ng/g of Re was detected in the chloroplast extract, and 3 ng/g of Re detected in the HEPES solution in which the chloroplasts were stored (Table 7). Regardless of the difficulty in isolating the chloroplast, less than 1% of the Re is present in the chloroplast relative to the host structure (non-fertile tips) which possesses ~150 ng/g.

- **Cytoplasmic proteins purification**

Cytoplasmic proteins (~48 µg) were purified from 2 grams of wet (i.e. 0.6 grams dry) *F. vesiculosus* non-fertile tips. Proteins possess sizes in excess of 5 kDa, and were only found in fractions 4 to 6 eluting (1 ml fractions were collected with a G25 column). No Re was observed in the elutions containing the proteins (Fig. 8). However, a total amount of ~200 ng

of Re was removed from the chromatography from elutions 10 to 14 with other unknown particles smaller than 5 kDa. Given the total volume of macroalgae used for the isolation of the protein (i.e. 0.6 grams of dry weight) this equates to a concentration of ~300 ng/g Re, as it is between the range of Re expected to be in the non-fertile tips, it can be stated that all Re from the tips structures was eluted.

Discussion

- Localization of Re within *F. vesiculosus* structures

The apical growth in the Phaeophyceae family is thought to occur by division of cells in cylindrical directions, with daughter cells generating a parenchymatous tissue construction [26]. Parenchyma tissue cells are capable of cell division if stimulated and can differentiate into specialized cells for photosynthesis, reproduction, growth and nutrient uptake. In Phaeophyceae, it is possible to distinguish five types of cells: epidermal cells, primary cortical cells, secondary cortical cells, medullary cells and hyphae [35]. The non-fertile tips are the apical meristems of *F. vesiculosus*, therefore it is composed of cells that can divide and differentiate, including photosynthetic cells. Although there is variability between the different macroalgae specimens collected, the relative levels of Re vary significantly within the macroalgae structures. There are significant differences (p -value < 0.05) between the amount of Re stored in the tips (~126 ng/g) versus Re stored in the remainder of the macroalgae (~74 ng/g) (Fig. 1). Furthermore, significant concentration of Re is found in the non-fertile tips which suggests a link between Re and the meristematic and photosynthetic specialized cells. More specifically, an average concentration of 313 ng/g of Re was found in the non-fertile tips, 122 ng/g Re in the fertile tips, 67 ng/g Re in the blades, 66 ng/g Re in the vesicles, 23 ng/g Re in the stipe and 21 ng/g Re in the holdfast. This suggests that Re is most likely stored in the photosynthetic structures and it is not involved in the reproductive

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

structures (receptacles). In herbaceous plants the distribution of Re is also higher in photosynthetic structures, with 86% of the plant Re reported to be at the leaves [36]. Bozhkov and Borisova [37] stated that, in plants, Re is accumulated in chlorophylls forming $Mg(ReO_4)_2$. However, no Re was found in the chloroplasts of *F. vesiculosus*, thus our study suggests that Re is not strongly bound by/to chlorophylls. The concentrations of Re in the chloroplast extraction and the HEPES solution where the chloroplasts were stored are 1 and 3 ng/g of Re, respectively (See Table 7). These concentrations are very low, much lower than the concentrations expected given the observed concentration on the tip structures (~100 ng/g).

It should be emphasised that the data in Table 1 shows that there is Re in all parts of *F. vesiculosus*, i.e. Re is not locally concentrated into a single structure, or a small number of structures, which means that Re is present in all cell types. In previous studies it was demonstrated that the cell surface is not the main accumulation site of Re in the brown macroalgae *Pelvetia fastigiata* [9]. As a result it would be expected that Re enters into the cell and remains in the cytoplasmic or a cell compartment. Moreover, Xiong *et al.* [15] made a macroalgae cell gel by chemically modifying brown macroalgae with sulphuric acid obtaining a gel of the macroalgae alginate and fucoidan matrix. The resulting gel had a high Re affinity and it was stated that amino acids were taking part in Re absorption, as it was observed in the IR (i.e. InfraRed) spectra that the intensity of the peaks corresponding to amino $-NH_2$ groups decreased after adsorption. Moreover, this fact was supported by removal of the amino acids of the gel (i.e. previously boiling the brown algae) which showed no adsorption of Re. Thus, this could mean that Re is not found in the cell wall in macroalgae, but interacts with cell membrane proteins or other molecules that contain $-NH_2$ groups in the cell, while not interacting with cytoplasmic proteins (see Fig. 8). As in this present study no disruption of the membranes was carried out it cannot be assumed that

membrane bound proteins were simultaneously extracted. Moreover, the method for protein detection used does not detect free amino acids, peptides (i.e. Glutathione, metallothioneins and phytochelatins) and proteins smaller than 3 kDa. Thus, it cannot be stated absolutely that Re is not protein bound because we cannot be sure to have isolated all the proteins, but it can be stated that it is not related to cytoplasmic proteins larger than 3 kDa or, if it is, the Re binding of the protein is sufficiently weak that the analytical protocol for protein isolation is capable of breaking any Re protein associated bond.

- Comparison of perrhenate compounds (HReO₄, NaReO₄, KReO₄ and NH₄ReO₄) uptake by cultured *F. vesiculosus* tips

An absorption study of Re onto organic polymers was undertaken by Kim *et al.* [17], who concluded that negatively charged perrhenate ions interacted with protonated amine groups in chitosan. The authors explain the adsorption by a combination of a Langmuir-Frendlich type mechanism and the electric diffuse double layer model. Our experiments show that all perrhenate salts have the same linear trendline (Fig. 3A) which strongly differs from perrhenate obtained from HReO₄ (Fig. 3B). This unexpected result highlights the importance of the chemical species of Re compound used for doping, which we further discuss below.

Perrhenate salts (NaReO₄, KReO₄ and NH₄ReO₄) are highly soluble in water with solubilities around 1.1 g/mL. It has been observed that cations are used as a symport for perrhenate uptake in animal cells [19]. Our results seem to show that H⁺ is the best counter ion for perrhenate uptake, therefore a greater uptake is observed when HReO₄ is used. Moreover, H⁺ could be increasing the conversion of -NH₂ groups of the macroalgae to -NH₃⁺, thus allowing perrhenate to bind. Therefore more polymers of glucosamine and amino groups in *F. vesiculosus* [15, 18] could be positively charged allowing more perrhenate binding, as it has been observed that perrhenate interacts strongly with polymers of glucosamine [17] and

1
2
3 383 amino groups [15]. Although the difference of such discrepancy cannot be resolved here,
4
5 384 uptake of ReO_4^- is observed no matter the what form of perrhenate compound used. The
6
7 385 mechanisms that control Re entry into the cells of macroalgae have not been identified. There
8
9
10 386 are many reports studying cation metal transporters, [38–40], but little is known about anion
11
12 387 transporters (pumps) of macroalgae. Phosphate, chloride, sulphate, nitrate and molybdate
13
14 388 transporters are all anion transporters reported in cells. Macroalgae could take up Re as
15
16 389 perrhenate instead of other substrates of these transporters. Other trace metals in seawater
17
18 390 exist, rather than as the free metal ion, as oxo-anions (e.g., perrhenate, chromate, vanadate,
19
20 391 molybdate, arsenate). The existing active transport pumps (e.g. sulphate, nitrate, phosphate)
21
22 392 could be taking up such metal oxo-anions or there could be metal-specific pumps [41]. It has
23
24 393 been observed that arsenate and phosphate have a common mechanism of uptake in bacteria
25
26 394 and yeast [42], but not in phytoplankton [43] and brown macroalgae [20], although high
27
28 395 concentrations of phosphate inhibit the uptake of arsenate. Nitrate could be also competing
29
30 396 with perrhenate, however this has only been observed for the mineral sodalite, and not in
31
32 397 living organisms [22].
33
34
35
36
37 398 The seasonal Re(VII) salt uptake variation of cultures (Table 6) suggest that perrhenate
38
39 399 uptake is biologically influenced. Riget *et al.* [44], observed that zinc obtained maximum
40
41 400 concentrations in macroalgae in March and a minimum in September, and it was similarly
42
43 401 observed, albeit less clearly, with lead and copper. Macroalgae growth is the most likely
44
45 402 cause for seasonal variations in metal uptake [44, 45]. Although our studies seem to support
46
47 403 this theory, a monthly perrhenate uptake research should be done in order to confirm it more
48
49 404 strongly and descipher if it is simply a dilution effect or if perrhenate has a real metabolic
50
51 405 role in the macroalgae. Here we did not perform any seasonal experiments using HReO_4 .
52
53
54
55
56 406 Our study also shows that when non-fertile thallus tips start dying they do not accumulate
57
58 407 more Re and start to degrade, thus Re is released to the media (Table 6; Fig. 4). Therefore,
59
60

less accumulation of Re in those cultured macroalgae tips that started dying is expected. This happened in the macroalgae tips cultured with 2000 and 7450 ng/g of HReO_4 in the seawater. In addition, it is worth emphasising that the more time the dying tips are left in the water, the more Re is released in the seawater by macroalgae (i.e. the less accumulation of Re). Thus, this explains the results obtained in Figure 4, where non-fertile thallus tips grown with a concentration of 2000 ng/g of HReO_4 accumulate less Re than the ones cultured with 7450 ng/g, because the firsts ones were cultured for 15 more days than the tips grown with 7450 ng/g of HReO_4 .

Therefore, a good linear correlation fit between HReO_4 doped in seawater and Re taken up by *F. vesiculosus* is observed up to 75 ng/g Re in seawater, but with higher concentrations (i.e. 1000, 2000 and 7450 ng/g) there is no linear correlation (Fig. 4 and 5) due to the probable metabolically inactivation of the tips. This indicates that the limit of uptake by the tips occurs when the tips are grown in a media of between 75 and 1000 ng/g of Re.

Phytoaccumulation (or phytoextraction) of metals by plants and algae is widely known [46], and refers to the concentration of metals from the environment into plant tissues. Plants absorb substances through the root, and then they transport and store these substances into the stems or leaves. There are two types of phytoextraction species: accumulator species and hyperaccumulator species. The main difference between those two types is stated in Rascio *et al.* [47]. Hyperaccumulator species are able to extract higher concentrations of metals and have a faster root-to-shoot transport system compared to non-hyperaccumulators species without showing phytotoxic effects. However, from the data obtained in this study it cannot be stated that *F. vesiculosus* is a hyperaccumulator species, because the thallus tips grown with the highest concentrations of ReO_4^- started to decrease in growth and die; although they were at concentrations not typical of any environmental setting.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

- **An understanding of Re uptake: active or passive**

Figure 6 and 7 show that Re uptake is not by simple diffusion, as it is observed that only living *F. vesiculosus* tips concentrate Re. Re levels in tips with high Re media concentration (7.45 ng/g) do not decrease when subsequently placed in media with lower Re concentrations: this suggests that the adsorption is not driven by simple equilibria. If Re was taken up by simple diffusion we would expect the same uptake of Re after boiling, freezing or drying the tips, as the membranes are not affected, and a direct correlation between the concentration of Re in the solution and in the macroalgae tips would be expected. Although Re could be taken up through passive mediated transport (facilitated diffusion) because after metabolically inactivating the macroalgae tips the transport proteins of the membranes are expected to be denatured (as happens when tips are boiled), thus no uptake is observed. However, this seems unlikely, due to the high Re uptake observed in living *F. vesiculosus* tips relative to the Re concentration in seawater. In addition, our results show that the uptake mechanism is syn-life, therefore Re is bioabsorbed. It can also be concluded that Re is not taken up by simple diffusion, at least for the perrhenate compounds used here. And, finally, Figure 6, shows that the uptake mechanism of the macroalgae is unidirectional, not a simple partition, as we observe that once living *F. vesiculosus* has accumulated Re, it does not release it back to the media.

- **Implications of bioaccumulation of Re**

Our results show little to no Re accumulation by metabolically inactivated *F. vesiculosus*, thus, if this is the case of macroalgae preserved in sediments as organic matter, using Re as a paleo redox may not strictly apply. However, we do suggest that once *F. vesiculosus* has died we may see release back to the water column as the macroalgae breaks down. Thus anoxia may be how the Re is stabilized, through prevention of macroalgae degradation.

Despite *F. vesiculosus* being a non-hyperaccumulator macroalgae, it is seen that until a limit, *F. vesiculosus* can accumulate up to 50000 ng/g when HReO_4 was present in the media, recovering the metal from the media. Thus, *F. vesiculosus* could be used as a source of phytomining of Re. Although differences in Re uptake are associated with the form of the perrhenate compounds, all ReO_4^- compounds used here permit the uptake of Re by *F. vesiculosus*. Moreover, as Re is also a Tc analogue [17], *F. vesiculosus* could be used for bioremediation of contaminated waters with Tc residues, as it has been found in ocean waters near to the Fukushima nuclear accident [49]. Tc is a radioactive metal mostly produced artificially during nuclear reactor fission product of uranium and plutonium.

Conclusions

The observation that macroalgae concentrates Re, an element with no known biological use, raises interesting questions. This study documents the first detailed examination of the relative proportions of Re in the structures of the macroalgae. The following conclusions are drawn from the present study:

- i. Re is not solely concentrated into a single macroalgae structure, all the cells possess Re. There is a distribution of Re that increases from the holdfast to the tips. Non-reproductive thallus tips exhibit the most Re accumulation, even more than reproductive thallus tips. As the only difference between them is the reproductive structures (receptacles), we can say that Re is not bound in the reproductive structures.
- ii. Our data shows that Re is bioadsorbed by *F. vesiculosus*, rather than bioaccumulated, and does not follow a simple diffusion uptake mechanism. The uptake is unidirectional, not a simple partition, however the data conclusively, *F. vesiculosus* uptakes and stores Re.

- 1
2
3 479 iii. Re recovery is observed from the seawater enriched with ReO_4^- , opening the possibility
4
5 480 of using *F. vesiculosus* as a source of phytomining.
6
7 481 iv. A differences in the uptake of Re between pherrenate salts and HReO_4 is observed,
8
9 482 however the cause has yet to be established.
10
11 483 v. The seasonal differences in Re uptake associated with pherrenate salts are a function of
12
13 484 *F. vesiculosus* growth.
14
15
16 485 vi. There is a limit on the uptake of Re in the cultured macroalage between 75 and 1000
17
18 486 ng/g of HReO_4 in the seawater media, and beyond that a deleterious effect is observed.
19
20
21 487 vii. Re is not accumulated in the cytoplasmic proteins or chloroplasts.
22
23

24 488 **References**

25
26
27 489 1. Koide M, Hodge VF, Yang JS, Stallard M, Goldberg EG, Calhoun J, Bertine KK. 1986
28
29 490 Some comparative marine chemistries of rhenium, gold, silver and molybdenum. *Appl.*
30
31 491 *Geochem.* **1**: 705–714. (doi: 10.1016/0883-2927(86)90092-2)
32
33
34 492 2. Anbar AD, Creaser RA, Papanastassiou DA, Wasserburg GJ. 1992 Rhenium in seawater:
35
36 493 Confirmation of generally conservative behavior. *Geochim. Cosmochim. Ac.* **56**: 4099–
37
38 494 4103. (doi: 10.1016/0016-7037(92)90021-A)
39
40 495 3. Helz GR, Dolor MK. 2012 What regulates rhenium deposition in euxinic basins? *Chem.*
41
42 496 *Geol.* **304–305**: 131–141. (doi: 10.1016/j.chemgeo.2012.02.011)
43
44
45 497 4. Miller CA, Peucker-Ehrebrink B, Walker BD, Marcantonio F. 2011 Re-assessing the
46
47 498 surface cycling of molybdenum and rhenium. *Geochim. Cosmochim. Ac.* **75**: 7146–7179.
48
49 499 (doi: 10.1016/j.gca.2011.09.005)
50
51 500 5. Selby D, Creaser RA. 2003 Re-Os geochronology of organic rich sediments: an
52
53 501 evaluation of organic matter analysis methods. *Chem. Geol.* **200**: 225–240. (doi:
54
55 502 10.1016/S0009-2541(03)00199-2)
56
57
58
59
60

- 503 6. Stein HJ. 2013 Dating and tracing the history of ore formation. *Treatise on Geochemistry*
504 **13**: 87–118. (doi: 10.1016/B978-0-08-095975-7.01104-9)
- 505 7. Mas JL, Tagami K, Uchida S. 2005 Rhenium measurements on North Atlantic seaweed
506 samples by ID-ICP-MS: An observation on the Re concentration factors. *J. Radioanal.*
507 *Nucl. Ch.* **265**: 361–365. (doi: 10.1007/s10967-005-0833-3)
- 508 8. Prouty NG, Roark EB, Koenig AE, Demopoulos AWJ, Batista FC, Kocar BD, Selby D,
509 McCarty MD, Mienis F, Ross SW. 2014 Deep-sea coral record of human impact on
510 watershed quality in the Mississippi River Basin. *Global Biogeochem. Cy.* **28**: 29–43.
511 (doi: 10.1002/2013GB004754)
- 512 9. Yang JS. 1991 High rhenium enrichment in brown algae: a biological sink of rhenium in
513 the sea? *Hydrobiologia* **211**: 165–170. (doi: 10.1007/BF00008532)
- 514 10. Davis TA, Volesky B, Mucci A. 2003 A review of the biochemistry of heavy metal
515 biosorption by brown algae. *Water Res.* **37**: 4311–4330. (doi: 10.1016/S0043-
516 1354(03)00293-8)
- 517 11. Chapman V, Chapman D. Seaweeds and Their Uses. 3rd ed. New York: 1980.
- 518 12. Lobban CS, Harrison PJ. Seaweed Ecology and Physiology. Cambridge University.
519 Cambridge: 1994.
- 520 13. Ragan MA, Smidsrod O, Larsen B. 1979 Chelation of divalent metal ions by brown algal
521 polyphenols. *Mar. Chem.* **7**: 265–271. (doi: 10.1016/0304-4203(79)90043-4)
- 522 14. Raize O, Argaman Y, Yannai S. 2004 Mechanisms of biosorption of different heavy
523 metals by brown marine macroalgae. *Biotechnol. Bioeng.* **87**: 451–458. (doi:
524 10.1002/bit.20136)
- 525 15. Xiong Y, Xu J, Shan W, Lou Z, Fang D, Zang S, Han G. 2013 A new approach for
526 rhenium(VII) recovery by using modified brown algae *Laminaria japonica* absorbent.
527 *Bioresource Technol.* **127**: 464–472. (doi: 10.1016/j.biortech.2012.09.099)

1
2
3 528 16. Meilián C, Kremer C, Suescun L, Mombrú A, Mariezcurrena R, Kremer E. 2000 Re(V)
4
5 529 complexes with amino acids based on the ‘3+2’ approach. *Inorg. Chim. Acta* **306**: 70–77.
6
7 530 (doi: 10.1016/S0020-1693(00)00151-1)
8
9
10 531 17. Kim E, Benedetti MF, Boulègue J. 2004 Removal of dissolved rhenium by sorption onto
11
12 532 organic polymers: study of rhenium as an analogue of radioactive technetium. *Water Res.*
13
14 533 **38**: 448–454. (doi: 10.1016/j.watres.2003.09.033)
15
16 534 18. Nishino T, Nishioka C, Ura H, Nagumo T. 1994 Isolation and partial characterisation of
17
18 535 a novel amino sugar-containing fucan sulfate from commercial *Fucus vesiculosus*
19
20 536 *fucoïdan*. *Carbohydr. Res.* **255**: 213–224. (doi: 10.1016/S0008-6215(00)90980-7)
21
22
23 537 19. Tagami K, Uchida S. 2005 A comparison of concentration ratios for technetium and
24
25 538 nutrient uptake by three plant species. *Chemosphere* **60**: 714–717. (doi:
26
27 539 10.1016/j.chemosphere.2005.03.087)
28
29
30 540 20. Klumpp DW. 1980 Characteristics of arsenic accumulation by the seaweeds *Fucus*
31
32 541 *spiralis* and *Ascophyllum nodosum*. *Mar. Biol.* **58**: 257–264. (doi: 10.1007/BF00390774)
33
34 542 21. Harvey BR, Williams KJ, Lovett MB, Ibbett RD. 1991 Determination of technetium-99
35
36 543 in environmental material with rhenium as a yield monitor. *J. Radioanal. Nucl. Ch.* **158**:
37
38 544 417–436. (doi: 10.1007/BF02047127)
39
40
41 545 22. Dickson JO, Harsh JB, Flury M, Lukens WW, Pierce EM. 2014 Competitive
42
43 546 incorporation of perrhenate and nitrate into a soladite *Environ. Sci. Technol.* **48**: 12851-
44
45 547 12857. (doi: 10.1021/es503156v)
46
47
48 548 23. Carlson L. 1991 Seasonal variation in growth, reproduction and nitrogen content of
49
50 549 *Fucus vesiculosus* in the Öresund, Southern Sweden. *Bot. Mar.* **34**: 447–453. (doi:
51
52 550 10.1515/botm.1991.34.5.447)
53
54
55 551 24. Strömberg T. 1977 Short-term effect of temperature upon the growth of intertidal
56
57 552 Fucales. *J. Exp. Mar. Biol. Ecol.* **29**: 181–195. (doi: 10.1016/0022-0981(77)90047-8)
58
59
60

- 553 25. White N. 2008 *Fucus vesiculosus*. Bladder wrack. Marine Life Information Network:
554 Biology and Sensitivity Key Information Sub-programme [on-line]. Plymouth: Marine
555 Biological Association of the United Kingdom. [cited 09/09/2015]. Available from:
556 <http://www.marlin.ac.uk/speciesinformation.php?speciesID=3348>
- 557 26. Graham LE, Wilcox LW. Algae. Prentice Hall. New Jersey; 2000.
- 558 27. Hiscock S. A Field Key to the British Brown Seaweeds (Phaeophyta). Field Studies
559 Council. Preston; 1991.
- 560 28. Gustow L, Rahman MM, Bartl K, Saborowski R, Bartsch I, Wiencke C. 2014 Ocean
561 acidification affects growth but not nutritional quality of the seaweed *Fucus vesiculosus*
562 (Phaeophyceae, Fucales). *J. Exp. Mar. Biol. Ecol.* **453**: 84–90. (doi:
563 10.1016/j.jembe.2014.01.005)
- 564 29. Gaines P. 1985-2014 Sample Preparation Guide [online]. Inorganic Ventures.
565 Christiansburg, Virginia USA. [cited 16/12/15]. Available from:
566 <http://www.inorganicventures.com/samples-containing-rhenium>
- 567 30. U.S. Geological Survey, 2007, Glossary--Bioaccumulation: U.S. Geological Survey,
568 accessed February 24, 2016
- 569 31. Popovic R, Colbow K, Vidaver W, Bruce D. 1983 Evolution of O₂ in Brown Algal
570 Chloroplasts. *Plant Physiol.* **73**: 889–892. (doi: 10.1104/pp.73.4.889)
- 571 32. de Boer E, Tromp MGM, Plat H, Krenn GE, Wever R. 1986 Vanadium(V) as an
572 essential element for haloperoxidase activity in marine brown algae: purification and
573 characterization of a vanadium(V)-containing bromoperoxidase from *Laminaria*
574 *saccharina*. *Bioch. Biophys. Acta* **872**: 104–115. (doi: 10.1016/0167-4838(86)90153-6)
- 575 33. GE Healthcare (2007) PD-10 Desalting Columns. Instructions 52-1308-00-88. Uppsala,
576 Sweeden.

1
2
3 577 https://www.gelifesciences.com/gehcls_images/GELS/Related%20Content/Files/131472
4
5 578 3116657/litdoc52130800BB_20110830191706.pdf
6
7
8 579 34. Cumming VM, Poulton SW, Rooney AD, Selby D. 2013 Anoxia in the terrestrial
9
10 580 environment during the late Mesoproterozoic. *Geology* **41**: 583–586. (doi:
11 581 10.1130/G34299.1)
12
13
14 582 35. Davy de Virville A, Feldmann J. IV International Symposium on Seaweed. Pergamon
15
16 583 Press. Biarritz; 1961
17
18
19 584 36. Bozhkov O, Christina T, Borisova L, Ernakov V, Ryabukhin V. 2006 Study of rhenium
20
21 585 accumulation in plants. *Trends in Inorganic Chemistry* **9**: 1–9.
22
23 586 37. Bozhkov O, Borisova L. 2002 Extraction and determination of trace amounts of
24
25 587 Rhenium in plants. *Int. J. Environ. An. Ch.* **83**: 135–141. (doi:
26
27 588 10.1080/0306731021000048627)
28
29
30 589 38. Cobbett CS, Hussain D, Haydon MJ. 2003 Structural and functional relationships
31
32 590 between type 1_B heavy metal-transporting P-type ATPases in *Arabidopsis*. *New Phytol.*
33
34 591 **159**: 315–321. (doi: 10.1046/j.1469-8137.2003.00785.x)
35
36
37 592 39. Mäser P, Thomine S, Schroeder JI, Ward JM, Hirschi K, Sze H, Talke IN, Amtmann A,
38
39 593 Maathuis FJM, Sanders D, Harper JF, Tchieu J, Gribskov M, Persans MW, Salt DE, Kim
40
41 594 SA, Guerinot ML. 2001 Phylogenetic Relationships within Cation Transporter Families
42
43 595 of *Arabidopsis*. *Plant Physiol.* **126**: 16461–667. (doi: 10.1104/pp.126.4.1646)
44
45 596 40. Williams LE, Pittman JK, Hall JL. 2000 Emerging mechanisms for heavy metal transport
46
47 597 in plants. *Biochim. Biophys. Acta* **1465**: 104–126. (doi: 10.1016/S0005-2736(00)00133-
48
49 598 4)
50
51
52 599 41. Dallinger R, Rainbow PS. Ecotoxicology of Metals in Invertebrates. CRC Press. Florida;
53
54 600 1993.
55
56
57
58
59
60

- 601 42. Rothstein A, Donovan K. 1963 Interactions of Arsenate with the Phosphate-Transporting
602 System of Yeast. *J. Gen. Physiol.* **46**: 1075–1085. (doi: 10.1085/jgp.46.5.1075)
- 603 43. Andrae MO, Klumpp DW. 1979 Biosynthesis and release of organoarsenic compounds
604 by marine algae. *Environ. Sci. Technol.* **13**: 738–741. (doi: 10.1021/es60154a001)
- 605 44. Riget F, Johansen P, Asmund G. 1995 Natural Seasonal Variation of Vanadium, Copper,
606 Lead and Zinc in Brown Seaweed (*Fucus vesiculosus*). *Mar. Pollut. Bull.* **30**: 409–413.
607 (doi: 10.1016/0025-326X(95)99847-W)
- 608 45. Fuge R, James KH. 1973 Trace metal concentrations in brown seaweeds, Cardigan Bay,
609 Wales. *Mar. Chem.* **1**: 281–293. (doi: 10.1016/0304-4203(73)90018-2)
- 610 46. Lasat MM. 2002 Phytoextraction of Toxic Metals: A Review of Biological Mechanisms.
611 *J. Environ. Qual.* **31**: 109–120. (doi: 10.2134/jeq2002.1090)
- 612 47. Rascio N, Navarri-Izzo F. 2011 Heavy metal hyperaccumulating plants: How and why
613 do they do it? And what makes them so interesting? *Plant Sci.* **180**: 169–181. (doi:
614 10.1016/j.plantsci.2010.08.016)
- 615 48. Steinhauser G. 2014 Fukushima's forgotten radionuclides: a review of the understudied
616 radioactive emissions. *Environ. Sci. Technol.* **48**: 4649–4663. (doi: 10.1021/es405654c)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

618 Table 1. Re abundance for *F. vesiculosus* structures analysed with Thermo Scientific X-
619 Series ICP-MS isotope dilution methodology.

Sample	Re (ng/g)	2σ (±)
Macroalgae 1		
Control	69.8	0.1
Tips 1	163.4	0.1
Leaves	28.4	0.1
Stipe	23.0	0.2
Holdfast	21.0	0.2
Blades	67.3	0.1
Veins	33.8	0.1
Blades without veins	65.8	0.1
Macroalgae 2		
Fertile tips	117.4	< 0.1
Non-Fertile tips	383.2	< 0.1
Tips	76.0	0.1
Control	51.0	0.1
Macroalgae 3		
Fertile tips	145.0	< 0.1
Non-Fertile tips	363.2	< 0.1
Tips	144.1	< 0.1
Control	103.4	0.1
Macroalgae 4		
Fertile tips	106.4	0.1
Non-Fertile tips	273.5	< 0.1
Tips	158.5	0.1
Control	61.0	0.1
Macroalgae 5		
Fertile tips	120.7	0.1
Non-Fertile tips	229.1	< 0.1
Tips	147.2	0.1
Control	84.3	0.1
Macroalgae 6		
Non-Fertile tips	382.5	< 0.1
Fertile tips	129.5	0.1
Tips	105.1	0.1
Macroalgae 7		
Control *	64.0	0.7
Tips *	138.0	0.7
Blades *	56.8	0.3
Stipe *	22.5	0.2
Holdfast *	21.6	0.2
Blades2 *	58.9	0.4

620 (*) samples analysed with Thermo Scientific Triton Mass Spectrometer.

Table 2. Re concentrations of the media utilized for Re uptake experiments for boiled (2 h and 5 min) and dried and freezed with liquid nitrogen *F. vesiculosus* tips. Re abundances determined with Thermo Scientific X-Series ICP-MS isotope calibration methodology.

Non-reproductive thallus tips treatment	Re (ng/g) doped in sea-water media previously	Re (ng/g) in seawater media afterwards	2 σ (\pm)
Boiled			
2 h	7.5	7.1	0.0
5 min	7.5	7.1	0.1
Dried 72 h	7.5	2.6	0.0
Freezed with N ₂ liquid	7.5	6.6	0.0
Non-treated Macroalgae (control)	7.5	0.3	0.0

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

628 Table 3. Re concentrations of the boiled (2 h and 5 min) and dried and freezed with liquid
629 nitrogen *F. vesiculosus* tips following Re uptake experiments. Re abundances determined
630 with Thermo Scientific X-Series ICP-MS isotope calibration methodology.

631

Non-reproductive thallus tips treatment	Re (ng/g) doped in seawater media	Re (ng/g) uptaken by <i>F. vesiculosus</i>	2σ (±)
Boiled			
2 h	7.5	36.2	0.1
2 h	0.0075	1.1	1.0
2 h	0.0	0.5	1.0
5 min	7.5	20.9	< 0.1
Dried 72 h	7.5	24.1	< 0.1
Freezed with N ₂ liquid	7.5	20.0	< 0.1

632
633
634

Table 4. Re concentration of Macroalgae tips cultured under the different concentrations of HReO₄ in the media. Re abundances determined with Thermo Scientific X-Series ICP-MS with isotope calibration methodology.

Replicate number	HReO ₄ (ng/g) Seawater	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
1	0.0075	187.0	0.4	168.2	9.5
2	0.0075	149.4	0.2		
1	0.07	549.6	0.2	415.4	50.6
2	0.07	391.0	0.1		
3	0.07	305.7	1.0		
1	0.4	995.2	16.0	1275.6	135.2
2	0.4	1190.0	1.3		
3	0.4	1641.7	52.0		
1	0.8	1668.1	0.3	1769.6	84.4
2	0.8	2007.3	3.0		
3	0.8	1633.3	2.4		
1	3.7	8575.0	18.1	9218.6	455.1
2	3.7	10505.9	2.9		
3	3.7	8575.0	12.8		
1	7.5	15961.8	37.9	16208.7	90.1
2	7.5	16387.0	5.0		
3	7.5	16277.3	50.2		
1	20.0	48738.7	69.0	48007.2	2009.2
2	20.0	52521.9	74.0		
3	20.0	42760.9	68.0		
1	75.0	51477.0	72.0	63283.4	5718.7
2	75.0	59611.8	16.5		
3	75.0	78761.5	99.0		
1	1000.0	53009.5	45.0	55588.2	2188.9
2	1000.0	61752.1	85.5		
3	1000.0	52003.1	99.5		
1	2000.0	23488.8	4.0	22472.5	512.0
2	2000.0	21070.8	26.5		
3	2000.0	22857.8	16.0		
1	7450.0	33061.0	50.0	33061	

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

642 Table 5. Re concentration of macroalgae tips cultured under the different concentrations of
643 Re(VII) salts and HReO₄ in the media. Re abundances determined with Thermo Scientific X-
644 Series ICP-MS with isotope calibration methodology.

645

Replicate number	NaReO ₄ (ng/g) Seawater (March)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	206.3	0.2	219.6	6.6
3	0.074	232.9	0.5		
2	0.373	624.5	0.8	629.5	2.5
3	0.373	634.5	1.0		
2	0.745	986.7	2.3	1033.6	23.4
3	0.745	1080.4	2.1		
2	7.450	8421.4	6.3	8064.2	178.6
3	7.450	7706.9	11.5		
Replicate numeber	NaReO ₄ (ng/g) Seawater (May)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.0074	95.3	< 0.1	86.1	4.6
3	0.0074	76.9	< 0.1		
2	0.074	175.0	< 0.1	132.9	21.0
3	0.074	90.9	< 0.1		
2	0.373	214.3	0.1	200.3	7.0
3	0.373	186.4	0.1		
2	0.745	227.9	0.3	225.7	1.1
3	0.745	223.5	0.2		
2	7.450	1268.0	1.1	1203.9	32.0
3	7.450	1139.9	1.7		
Replicate number	NH ₄ ReO ₄ (ng/g) Seawater (May)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	230.6	< 0.1	226.1	2.2
3	0.074	221.6	< 0.1		
2	0.373	128.6	< 0.1	129.4	9.4

3	0.373	130.1	< 0.1		
2	0.745	283.6	< 0.1		
3	0.745	254.3	0.1	268.9	7.3
2	7.450	1244.6	0.3		
3	7.450	1171.6	2.1	1208.1	18.2
Replicate number	KReO ₄ (ng/g) Seawater (May)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	88.0	0.1		
3	0.074	95.9	0.1	91.9	7.0
2	0.373	143.6	< 0.1		
3	0.373	133.2	0.1	138.4	2.6
2	0.745	166.5	< 0.1		
3	0.745	185.8	0.3	176.1	4.8
2	7.450	1260.3	0.5		
3	7.450	1242.2	0.6	1251.1	4.4
Replicate number	NH ₄ ReO ₄ (ppb) Seawater (June)	Re (ppb) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	81.0	0.2		
1	0.074	83.7	< 0.1	82.3	0.7
2	0.745	125.4	0.2		
1	0.745	133.0	< 0.1	129.2	1.9
2	7.450	689.2	3.3		
1	7.450	776.4	0.2	732.8	21.8
Replicate number	KReO ₄ (ng/g) Seawater (June)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	51.9	0.1		
1	0.074	64.6	< 0.1	58.3	3.2
2	0.745	233.8	0.6		
1	0.745	242.6	1.0	272.4	2.2
2	7.450	587.0	0.4		
1	7.450	544.4	< 0.1	564.9	10.7

Replicate number	HReO ₄ (ng/g) Seawater (June)	Re (ng/g) uptake by <i>F. vesiculosus</i>	2σ (±)	Replicates average	SD (±)
2	0.074	125.6	< 0.1	128.6	1.5
1	0.074	131.8	< 0.1		
2	0.745	733.79	0.2	722.5	5.6
1	0.745	711.3	41.0		
2	7.450	5924.3	33.5	6741.4	408.6
1	7.450	7558.6	56.5		

646

647

Table 6. Seasonal uptake percentage variation of Re(VII) salts (i.e. NH_4ReO_4 , KReO_4 and NaReO_4) cultures done in 2015 versus uptake rate of HReO_4 cultures performed in June 2014 and 2015.

	Re(VII) salts				HReO ₄	
	February 2015	March 2015	May 2015	June 2015	June 2014	June 2015
Number of media changes	5	5	7	4	5	4
Total ReO ₄ (ng) in seawater [doped ng × num. of media changes]	12500	12500	17500	10000	9300	7440
Possible Re (ng/g) accumulation by <i>F. v.</i> *	~25000	~2500 0	~3500 0	~2000 0	~18600	~1488 0
Real Re (ng/g) accumulation by <i>F.v.</i>	~1700	~8000	~1200	~800	~9300	~7400
% Uptake [Real / Possible accumulation]	6,80%	32,00 %	3,40%	4,00%	50,00%	49,70 %

* Total Re in seawater / average dry weight of macroalgae tips (0.5 g)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

653 Table 7. Concentration of Re (ng/g) in Chloroplasts and in HEPES solution where
654 chloroplasts were stored.

Sample	Re concentration (ng/g)
Chloroplast pellet	~ 1
HEPES solution	~ 3

655

Fig. 1 Average (2-5 samples) concentration of Rhenium (ng/g) in the different structures of *F. vesiculosus*. Round marker symbolizes Re abundance in each particular structure and square marker symbolizes Re abundance of a mixture of all the structures (control). All the samples had a reproducibility of < 5% RSD, in some cases, graph symbol size is greater than uncertainties. The concentrations shown are in dry mass, and although the concentration of each structure might change when wet mass, the differences of Re concentration are greater than the differences in water loss.

Fig. 2 Culture representation of non-reproductive *F. vesiculosus* thallus tips. 21 tips of each *F. vesiculosus* specimen were cut and a tip from each specimen was displaced into one of the 21 jars (A). Two meshes were put inside each jar ending up with three levels that store three non-fertile tips each (B). C) Real culture jar picture.

Fig. 3 A) Rhenium (ng/g) accumulation in *F. vesiculosus* under different Re(VII) salts concentrations. Cultures made with NH_4ReO_4 represented with a round marker, KReO_4 shown in square marker and NaReO_4 in triangle marker. **B)** Rhenium (ng/g) accumulation in *F. vesiculosus* under different Re(VII) salts (round marker) and HReO_4 (square marker) plotted in logarithmic scale. All the samples had a reproducibility of <5% RSD, in some cases, graph symbol size is greater than uncertainties.


Fig. 4 Rhenium (ng/g) accumulation in *F. vesiculosus* under different HReO_4 doped seawater concentrations. It follows a logarithmic trend line. All the samples had a reproducibility of < 5% RSD, in some cases, graph symbol size is greater than uncertainties.

Fig. 5 Rhenium (ng/g) accumulation in *F. vesiculosus* under different HReO_4 doped seawater concentrations. All the samples had a reproducibility of <5% RSD, in some cases, graph symbol size is greater than uncertainties.

Fig. 6 Re (ng/g) accumulation in *F. vesiculosus* under changing concentrations of Re(VII) salts in the media. Day 1 to 3 Re concentration of 7.45 ng/g, from day 3 to 6; 0.075 ng/g and

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

681 from day 6 to 9; 0.0075 ng/g. Day 0 measure is the background concentration of Re found in
682 the seaweed cultured. All the samples had a reproducibility of <5% RSD.
683 **Fig. 7** Accumulation of ReO_4^- in *F. vesiculosus* under different treatments (previously heated
684 at 100 °C for 5 min, liquid nitrogen freezed and 30 °C dried) and 7.45 ng/g HReO_4 media
685 concentration. All the samples had a reproducibility of < 5% RSD.
686 **Fig. 8 A)** Concentration of proteins ($\mu\text{g/mL}$) in each elution (i.e. fraction eluted,
687 corresponding to 1 mL). There are two protein peaks in elution 6 and 8-9. **B)** Concentration
688 of Rhenium (ng/g) in each elution. The peak is in the elution 12.



1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

